Sum rule of the correlation function

Radosław Maj*
Institute of Physics, Świętokrzyska Academy
ul. Świętokrzyska 15, PL - 25-406 Kielce, Poland

Stanisław Mrówczyński†
Sołtan Institute for Nuclear Studies
ul. Hoża 69, PL - 00-681 Warsaw, Poland
and Institute of Physics, Świętokrzyska Academy
ul. Świętokrzyska 15, PL - 25-406 Kielce, Poland
(5-th December 2004)

arXiv:nucl-th/0409061v2 28 Feb 2005

We discuss a sum rule satisfied by the correlation function of two particles with small relative momenta. The sum rule, which results from the completeness condition of the quantum states of the two particles, is derived and checked to see how it works in practice. The sum rule is shown to be trivially satisfied by free particle pairs, and then three different systems of interacting particles are considered. We discuss neutron and proton pairs in the $s-$wave approximation and the case of the so-called hard spheres with the phase shifts taken into account up to $l = 4$. Finally, the Coulomb system of two charged particles is analyzed.

PACS number: 25.75.Gz

I. INTRODUCTION

The correlation functions of two identical or non-identical particles with 'small' relative momenta have been extensively studied in nuclear collisions for bombarding energies from tens of MeV [1] to hundreds of GeV [2]. These functions provide information about space-time characteristics of particle sources in the collisions. As shown by one of us [3], the correlation function integrated over particle relative momentum satisfies a simple and exact relation due to the completeness of the particle quantum states. The relation can be used to get a particle phase-space density, following the method [4,5], with no need to extract the Coulomb interaction. The sum-rule offers a constraint for the procedure of imaging [6,7] which inverts the correlation function and provides the source function. The sum-rule is also helpful to handle the correlation functions of exotic systems like $\bar{p}\Lambda$ [8] when the interparticle correlation is poorly known.

The aim of this paper is to discuss the sum rule in detail, to prove or disprove its usefulness in the experimental studies. Therefore, we first derive the sum rule and show that it is trivially satisfied by free particles. Then, we analyze the correlations in the neutron-proton (n-p) system where there are both the two-particle scattering states and a bound state *i.e.* a deuteron. We prove that in spite of the attractive interaction the n-p correlation can be negative due to the deuteron formation. Although some qualitative features dictated by the sum rule are certainly seen, the calculated correlation function does not satisfy the sum rule. This is not surprising as not only the momenta of order $1/R$, where R is the source size, but also larger momenta contribute to the sum-rule integral. Consequently, the $s-$wave approximation, which is used compute the n-p correlation function, is insufficient.

To see how important are the higher l phase shifts, we discuss a system of the so-called hard spheres and compute the correlation function of identical and non-identical particles, taking into account the phase shifts up to $l = 4$. Paradoxically, the higher l contributions do not improve the situation but make it even worse.

Finally, we discuss the correlations due to the Coulomb interaction. This is the case of particular interest as one usually measures the correlations functions of charge particles which experience the Coulomb interaction. And the integral, which is controlled by the sum rule, is used to determine the particle phase-space density [4,5]. In the case of Coulomb interactions, the exact wave functions are known, and consequently, the exact correlation functions can

*Electronic address: radcypmaj@poczta.onet.pl
†Electronic address: mrow@fuw.edu.pl

be computed. Unfortunately, as we discuss in detail, the integral of interest appears to be divergent, and the sum rule does not hold for this most important case.

II. PRELIMINARIES

To avoid unnecessary complications, our considerations are non-relativistic and we start with the formula repeatedly discussed in the literature which expresses the correlation function $R(\mathbf{q})$ of two particles with the relative momentum $\mathbf{q}$ as

$$R(\mathbf{q}) = \int d^3r\, \mathcal{D}_r(\mathbf{r})\, |\phi_{\mathbf{q}}(\mathbf{r})|^2 \,, \tag{1}$$

where $\phi_{\mathbf{q}}(\mathbf{r})$ is the wave function of relative motion of the two particles and $\mathcal{D}_r(\mathbf{r})$ is the *effective* source function defined through the probability density $D_r(\mathbf{r},t)$ to emit the two particles at the relative distance $\mathbf{r}$ and the time difference t as

$$\mathcal{D}_r(\mathbf{r}) = \int dt\, D_r(\mathbf{r}-\mathbf{v}t,t) \,, \tag{2}$$

with $\mathbf{v}$ being the particle pair velocity with respect to the source; the relative source function $D_r(\mathbf{r},t)$ is given by the single-particle source function $D(\mathbf{r},t)$, which describes the space-time emission points of a single particle, in the following way

$$D_r(\mathbf{r},t) = \int d^3R dT\, D(\mathbf{R}-\frac{1}{2}\mathbf{r}, T-\frac{1}{2}t)\, D(\mathbf{R}+\frac{1}{2}\mathbf{r}, T+\frac{1}{2}t) \,.$$

We note that $\mathcal{D}_r(\mathbf{r})$, $D_r(\mathbf{r},t)$ and $D(\mathbf{r},t)$ are all normalized as

$$\int d^3r\, \mathcal{D}_r(\mathbf{r}) = \int d^3r\, dt\, D_r(\mathbf{r},t) = \int d^3r\, dt\, D_r(\mathbf{r},t) = 1 \,. \tag{3}$$

We also observe that the spherically symmetric single-particle source function $D(\mathbf{r},t)$ provides, in general, the effective $\mathcal{D}_r(\mathbf{r})$ which is elongated along the velocity $\mathbf{v}$. To simplify calculations, we, however, assume here that the source function $\mathcal{D}_r(\mathbf{r})$ is spherically symmetric. Such an assumption makes sense when the single-particle source function $D(\mathbf{r},t)$ is spherically symmetric and particles are emitted instantaneously *i.e.* $D(\mathbf{r},t) = D(|\mathbf{r}|)\,\delta(t-t_0)$. Then,

$$\mathcal{D}_r(\mathbf{r}) = \int d^3R\, D(\mathbf{R}-\frac{1}{2}\mathbf{r})\, D(\mathbf{R}+\frac{1}{2}\mathbf{r}) \,.$$

The single-particle source function is often chosen in the Gaussian form

$$D(\mathbf{r}) = \frac{1}{(2\pi r_0^2)^{3/2}} \exp\Big(-\frac{\mathbf{r}^2}{2r_0^2}\Big) \,.$$

It gives the mean radius squared of a source equal $\langle \mathbf{r}^2 \rangle = 3r_0^2$, and it leads to the effective relative source function as

$$\mathcal{D}_r(\mathbf{r}) = \frac{1}{(4\pi r_0^2)^{3/2}} \exp\Big(-\frac{\mathbf{r}^2}{4r_0^2}\Big) \,. \tag{4}$$

III. SUM RULE

Let us consider the correlation function integrated over the relative momentum. Since $R(\mathbf{q}) \to 1$ when $\mathbf{q} \to \infty$, we rather discuss the integral of $R(\mathbf{q})-1$. Using Eq. (1) and taking into account the normalization condition of $\mathcal{D}_r(\mathbf{r})$ (3), one finds, after changing order of the $\mathbf{r}-$ and $\mathbf{q}-$integration, the expression

$$\int \frac{d^3q}{(2\pi)^3} \Big(R(\mathbf{q})-1\Big) = \int d^3r\, \mathcal{D}_r(\mathbf{r}) \int \frac{d^3q}{(2\pi)^3} \Big(|\phi_{\mathbf{q}}(\mathbf{r})|^2 - 1\Big) \,. \tag{5}$$

It appears that the integral over $\mathbf{q}$ in the r.h.s. of Eq. (5) is determined by the quantum-mechanical completeness condition. Indeed, the wave functions satisfy the well-known closure relation

$$\int \frac{d^3q}{(2\pi)^3}\ \phi_{\mathbf{q}}(\mathbf{r})\phi^*_{\mathbf{q}}(\mathbf{r}') + \sum_\alpha \phi_\alpha(\mathbf{r})\phi^*_\alpha(\mathbf{r}') = \delta^{(3)}(\mathbf{r}-\mathbf{r}') \pm \delta^{(3)}(\mathbf{r}+\mathbf{r}')\ , \tag{6}$$

where ϕ_α represents a possible bound state of the two particles of interest. When the particles are not identical the second term in the r.h.s. of Eq. (6) is not present. This term guarantees the right symmetry for both sides of the equation for the case of identical particles. The upper sign is for bosons while the lower one for fermions. The wave function of identical bosons (fermions) $\phi_{\mathbf{q}}(\mathbf{r})$ is (anti-)symmetric when $\mathbf{r} \to -\mathbf{r}$, and the r.h.s of Eq. (6) is indeed (anti-)symmetric when $\mathbf{r} \to -\mathbf{r}$ or $\mathbf{r}' \to -\mathbf{r}'$. If the particles of interest carry spin, the summation over spin degrees of freedom in the l.h.s. of Eq. (6) is implied. When the integral representation of $\delta^{(3)}(\mathbf{r}-\mathbf{r}')$ is used, the relation (5) can be rewritten as

$$\int \frac{d^3q}{(2\pi)^3}\ \left(\phi_{\mathbf{q}}(\mathbf{r})\phi^*_{\mathbf{q}}(\mathbf{r}') - e^{i\mathbf{q}(\mathbf{r}-\mathbf{r}')}\right) + \sum_\alpha \phi_\alpha(\mathbf{r})\phi^*_\alpha(\mathbf{r}') = \pm\delta^{(3)}(\mathbf{r}+\mathbf{r}')\ .$$

Now, we take the limit $\mathbf{r}' \to \mathbf{r}$ and get the relation

$$\int \frac{d^3q}{(2\pi)^3}\ \left(|\phi_{\mathbf{q}}(\mathbf{r})|^2 - 1\right) = \pm\ \delta^{(3)}(2\mathbf{r}) - \sum_\alpha |\phi_\alpha(\mathbf{r})|^2\ . \tag{7}$$

When Eq. (7) is substituted into Eq. (5), one finds the desired sum rule

$$\int d^3q\ \left(R(\mathbf{q}) - 1\right) = \pm\pi^3\ \mathcal{D}_r(0) - \sum_\alpha A_\alpha\ . \tag{8}$$

where A_α is the formation rate of a bound state α

$$A_\alpha = (2\pi)^3 \int d^3r\ \mathcal{D}_r(\mathbf{r})|\phi_\alpha(\mathbf{r})|^2\ . \tag{9}$$

A_α relates the cross section to produce the bound state α with the momentum $\mathbf{P}$ to the cross section to produce the two particles with the momenta $\mathbf{P}/2$ as

$$\frac{d\sigma^\alpha}{d\mathbf{P}} = A_\alpha\ \frac{d\ \widetilde{\sigma}}{d(\mathbf{P}/2)d(\mathbf{P}/2)}\ .$$

The tilde means that the short range correlations are removed from the two-particle cross section which is usually taken as a product of the single-particle cross sections.

The completeness condition is, obviously, valid for any inter-particle interaction. It is also valid when the pair of particles interact with the time-independent external field, *e.g.* the Coulomb field generated by the particle source. Thus, the sum rule (8) holds under very general conditions as long as the basic formula (1) is justified, in particular as long as the source function $\mathcal{D}_r(\mathbf{r})$ is $\mathbf{q}-$independent and spin independent. The validity of these assumptions can be only tested within a microscopic model of nucleus-nucleus collision which properly describes the quantum particle correlations and bound state formation.

There are several potential applications of the relation (8). As already mentioned, the integral of the correlation function in the l.h.s. of Eq. (8) is used to determine a particle phase-space density. The method [4,5] assumes that the correlation function represents non-interacting particles. Consequently, before computing the integral, the correlation function is corrected in such a way that the Coulomb interaction is removed. However, the sum rule (8) shows that the integral of the correlation function is independent of the inter-particle interaction. Therefore, there is no need to extract the Coulomb interaction. We note that this procedure is rather model dependent.

To get information about the source function $D(\mathbf{r},t)$, one usually parameterizes the function, computes the correlation function and compares it with experimental data. The method of imaging [6,7] provides the source function directly, inverting the functional $R[\mathcal{D}_r]$ (1) with the experimental correlation function as an input. As seen, the relation (8), which gives $\mathcal{D}_r(0)$, can be treated as a useful constraint of the imaging method.

The sum-rule is also helpful to better understand the correlation functions. In particular, the sum rule shows that the correlation function can be negative in spite of an attractive inter-particle interaction. This happens when the particles form bound states represented by the second term in the r.h.s. of Eq. (8) or the interaction is strongly absorptive as in the case of antiproton-lambda system which annihilates into mesons. Then, the scattering states of the $\bar{p}\Lambda$ do not give a complete set of quantum states of the system and the integral from the l.h.s. of Eq. (8) has to be negative. The recently measured correlation function of $\bar{p}$ and Λ [8] is indeed negative at small relative momenta.

IV. FREE PARTICLES

In the case of non-interacting particles, the correlation function differs from unity only for identical particles. Then, the wave function, which enters the correlation function (1), is an (anti-)symmetrized plane wave

$$\phi_{\mathbf{q}}(\mathbf{r}) = \frac{1}{\sqrt{2}}\Big(e^{i\mathbf{qr}} \pm e^{-i\mathbf{qr}}\Big) \,,$$

with the upper sign for bosons and lower for fermions. Then, the integration over $\mathbf{q}$ in Eq. (5) can be explicitly performed without reference to the completeness condition (7), and one finds

$$\int d^3q \,\Big(R(\mathbf{q}) - 1\Big) = \pm\pi^3 \,\mathcal{D}_r(0) \,. \tag{10}$$

It this way the sum rule (10) was found in [9], see also [4].

Although the sum rule (8) assumes the integration up to the infinite momentum, one expects that the integral in Eq. (8) saturates at sufficiently large q. To discuss the problem quantitatively, we define the function

$$S(q_{\rm max}) = 4\pi \int_0^{q_{\rm max}} dq\, q^2 (R(q) - 1) \,. \tag{11}$$

As already mentioned, the source function is assumed to be spherically symmetric, and consequently the correlation function depends only on $q \equiv |\mathbf{q}|$. Thus, the angular integration is trivially performed.

For the Gaussian effective source (4), when $\pi^3\mathcal{D}_r(0) = (\sqrt{\pi}/2r_0)^3$, the free functions $R(q)$ and $S(q_{\rm max})$ equal

$$R(q) = 1 \pm e^{-4r_0^2 q^2} \,,$$

$$S(q_{\rm max}) = \pm\left(\frac{\sqrt{\pi}}{2r_0}\right)^3 \mathrm{E}_{2/3}\big((2r_0 q_{\rm max})^3\big) \,,$$

where

$$\mathrm{E}_n(x) \equiv \frac{1}{\Gamma(1+1/n)} \int_0^x dt\, e^{-t^n} \,,$$

and $\Gamma(z)$ is the Euler gamma function. Since for large x, the function $\mathrm{E}_n(x)$ can be expressed as, see *e.g.* [10],

$$\mathrm{E}_n(x) = 1 - \frac{1}{\Gamma(1+1/n)} \,\frac{e^{-x^n}}{n x^{n-1}} \left(1 + \mathcal{O}\Big(\frac{1}{x^n}\Big)\right) \,,$$

we have the approximation

$$S(q_{\rm max}) \approx \pm\left(\frac{\sqrt{\pi}}{2r_0}\right)^3 \Big(1 - \frac{4r_0 q_{\rm max}}{\sqrt{\pi}}\, e^{-4r_0^2\, q_{\rm max}^2}\Big) \,,$$

for $(2r_0\, q_{\rm max})^3 \gg 1$. As seen, the sum-rule integral (10) is saturated for $q_{\rm max}$ not much exceeding $1/2r_0$, and obviously, the sum rule is satisfied.

V. NEUTRON-PROTON SYSTEM

In this section, we discuss the interacting neutron-proton pair which is either in the spin singlet or triplet state. The nucleons, produced in high-energy nuclear collisions, are usually assumed to be unpolarized, and one considers the spin-averaged correlation function R which is a sum of the singlet and triplet correlation functions $R^{s,t}$ with the weight coefficients 1/4 and 3/4, respectively. Here, we consider, however, the singlet and the triplet correlation functions separately. Then, the sum rule (8) reads

$$\int d^3q \left(R^s(\mathbf{q}) - 1\right) = 0 \,, \tag{12}$$

$$\int d^3q \left(R^t(\mathbf{q}) - 1\right) = -A_d \,. \tag{13}$$

Following [11], we calculate the correlation functions $R^{s,t}$ assuming that the source radius is significantly larger than the n-p interaction range. Then, the wave function of the n-p pair (in a scattering state) can be approximated by its asymptotic form

$$\phi^{s,t}_{np}(\mathbf{r}) = e^{i\mathbf{qr}} + f^{s,t}(q)\frac{e^{iqr}}{r} \,, \tag{14}$$

where $f^{s,t}(\mathbf{q})$ is the scattering amplitude. It is chosen as

$$f^{s,t}(q) = \frac{-a^{s,t}}{1 - \frac{1}{2}d^{s,t}a^{s,t}q^2 + iqa^{s,t}} \,, \tag{15}$$

where $a^{s,t}$ ($d^{s,t}$) is the scattering length (effective range) of the n-p scattering; $a^s = -23.7$ fm, $d^s = 2.7$ fm and $a^t = 5.4$ fm, $d^t = 1.7$ fm [12]. The amplitude (15) takes into account only the $s-$wave scattering. This is justified as long as only 'small' relative momenta are considered.

Substituting the wave function (14) into the formula (1) with the source function (4), we get

$$R^{s,t}(q) = 1 + \mathrm{Re}\big[f^{s,t}(q)\big]\,\frac{1}{2r_0^2 q}\,e^{-4r_0^2q^2}\,\mathrm{erfi}(2r_0q) - \mathrm{Im}\big[f^{s,t}(q)\big]\,\frac{1}{2r_0^2 q}\big(1 - e^{-4r_0^2q^2}\big) + |f^{s,t}(q)|^2\frac{1}{2r_0^2} \,, \tag{16}$$

where

$$\mathrm{erfi}(x) \equiv \frac{2}{\sqrt{\pi}}\int_0^x dt e^{t^2} \,.$$

Since the source described by the formula (4) is spherically symmetric, the correlation function (16) does not depend on $\mathbf{q}$ but on q only.

In Figs. 1 and 2 we show, respectively, the singlet and triplet correlation functions computed for three values of the source size parameter r_0. As seen, the triplet correlation is negative in spite of the attractive neutron-proton interaction. This happens, in accordance with the sum rule (13), because the neutron and proton, which are close to each other in the phase-space, tend to exist in a bound not in a scattering state. And the n-p pairs, which form a deuteron, deplete the sample of n-p pairs and produce a dip of the correlation function at small relative momenta.

The deuteron formation rate, which enters the sum rule (13), is computed with the deuteron wave function in the Hulthén form

$$\phi_d(\mathbf{r}) = \Big(\frac{\alpha\beta(\alpha+\beta)}{2\pi(\alpha-\beta)^2}\Big)^{1/2}\;\frac{e^{-\alpha r} - e^{-\beta r}}{r} \,, \tag{17}$$

with $\alpha = 0.23$ fm^{-1} and $\beta = 1.61$ fm^{-1} [13]. Substituting the wave function (17) and the source function (4) into Eq. (9), we get

$$A_d = \frac{2\pi^2}{r_0^2}\,\frac{\alpha\beta(\alpha+\beta)}{(\alpha-\beta)^2}\big[K(2\alpha r_0) - 2K((\alpha+\beta)r_0) + K(2\beta r_0)\big] \,, \tag{18}$$

where

$$K(x) \equiv e^{x^2}\mathrm{erfc}(x) \,, \qquad \mathrm{erfc}(x) \equiv \frac{2}{\sqrt{\pi}}\int_x^\infty dt e^{-t^2} \,.$$

In Fig. 3 we present the deuteron formation rate (18) as function of the source size parameter r_0. As seen, A_d monotonously decreases when the source grows.

In Figs. 4 and 5 we display the function $S(q_{\mathrm{max}})$, defined by Eq. (11), found for the singlet and triplet correlation functions presented in Figs. 1 and 2, respectively. Although $S(q_{\mathrm{max}})$ saturates at large q_{max} for both the singlet and for the triplet correlation function, neither the sum rule (12) nor (13) is satisfied. The singlet $S(q_{\mathrm{max}})$ does not vanish at large q_{max}, as it should according to Eq. (12), while the triplet $S(q_{\mathrm{max}})$ is not negative, as required by

Eq. (13). However, comparing the numerical values of $S(q_{\max})$ to the corresponding deuteron formation rate, which sets the characteristic scale, one sees that with growing r_0, the sum rule is violated less and less dramatically. It is not surprising as the asymptotic form of the wave function (14) and the $s-$wave approximation become then more accurate. The formula (11) shows that due to the factor q^2 even small deviations of the correlation function from unity at large q generate sizable contribution to S. Therefore, it can be expected that higher partial waves have to be taken into account to satisfy the sum rule.

VI. HARD SPHERES

We consider here a pair of the so-called hard spheres interacting via the potential which is zero at distances larger than a and it is infinite otherwise; a is the sphere's diameter. As well known, see *e.g.* [14], scattering of the spheres is analytically tractable and the $l-$th phase shift δ_l is given by the equation

$$\mathrm{tg}\delta_l = \frac{j_l(qa)}{n_l(qa)} \,, \tag{19}$$

where $j_l(z)$ and $n_l(z)$ is the spherical Bessel and Neumann function, respectively. The scattering amplitude equals

$$f(\Theta) = \frac{1}{q}\sum_{l=0}^{\infty}(2l+1)\,P_l(\cos\Theta)\frac{\mathrm{ctg}\delta_l + i}{\mathrm{ctg}^2\delta_l + 1} \,, \tag{20}$$

where Θ is the scattering angle and $P_l(z)$ is the $l-$th Legendre polynomial. Substituting the amplitude (20) into the equation analogous to Eq. (14), we get the (asymptotic) wave function. Using this function and the Gaussian source (4), we get the correlation function of non-identical hard spheres as

$$\begin{aligned} R(q) = 1 + \frac{1}{\sqrt{\pi}\,r_0^3 q}\sum_l (2l+1)\,\frac{j_l(qa)}{n_l^2(qa)+j_l^2(qa)} \\ \times\Bigg[\Big(n_l(qa)\cos\Big(\frac{\pi l}{2}\Big) + j_l(qa)\sin\Big(\frac{\pi l}{2}\Big)\Big)\int_0^\infty drr\,e^{-\frac{r^2}{4r_0^2}}\cos(qr)\,j_l(qr) \\ +\Big(n_l(qa)\sin\Big(\frac{\pi l}{2}\Big) - j_l(qa)\cos\Big(\frac{\pi l}{2}\Big)\Big)\int_0^\infty drr\,e^{-\frac{r^2}{4r_0^2}}\sin(qr)\,j_l(qr)\Bigg] \\ + \frac{1}{2r_0^2q^2}\sum_l (2l+1)\,\frac{j_l^2(qa)}{n_l^2(qa)+j_l^2(qa)} \,. \end{aligned} \tag{21}$$

For identical hard spheres, the wave function should be (anti-)symmetrized as $\frac{1}{\sqrt{2}}\big(\phi_{\mathbf{q}}(\mathbf{r})\pm\phi_{\mathbf{q}}(-\mathbf{r})\big)$, and the correlation function equals

$$\begin{aligned} R(q) = 1 \pm e^{-4r_0^2q^2} + \frac{1}{\sqrt{\pi}\,r_0^3 q}\sum_l (2l+1)\,\frac{j_l(qa)}{n_l^2(qa)+j_l^2(qa)}\,[1\pm(-1)^l] \\ \times\Bigg[\Big(n_l(qa)\cos\Big(\frac{\pi l}{2}\Big) - j_l(qa)\sin\Big(\frac{\pi l}{2}\Big)\Big)\int_0^\infty drr\,e^{-\frac{r^2}{4r_0^2}}\cos(qr)\,j_l(qr) \\ -\Big(n_l(qa)\sin\Big(\frac{\pi l}{2}\Big) + j_l(qa)\cos\Big(\frac{\pi l}{2}\Big)\Big)\int_0^\infty drr\,e^{-\frac{r^2}{4r_0^2}}\sin(qr)\,j_l(qr)\Bigg] \\ + \frac{1}{2r_0^2q^2}\sum_l (2l+1)\,\frac{j_l^2(qa)}{n_l^2(qa)+j_l^2(qa)}\,[1\pm(-1)^l] \,. \end{aligned} \tag{22}$$

We note that there is a specific asymmetry of the signs of the corresponding terms in Eq. (21) and in Eq. (22). This happens because $((-1)^l\pm 1)/2 = \mp(-1)^l$.

In Figs. 6-9 we show the correlation functions for non-identical spheres (21) and identical bosonic ones (22). The functions are computed numerically for the sphere diameter $a = 1$ fm and two values of the source size parameter $r_0 = 4$ fm and $r_0 = 6$ fm. We have considered rather large sources as the wave function, which is used to compute the correlation function, is of the asymptotic form (14). This requires $r_0 \gg a$. The functions $S(q_{\rm max})$ defined by Eq. (11), which correspond to the correlation functions shown in Figs. 6-9, are presented in Figs. 10-13. There are in Figs. 6-13 three families of curves representing: the $s-$wave approximation, the sum of partial waves with $l = 0$, 1, 2, and the sum of $l = 0$, 1, 2, 3, 4. As seen, going beyond the $s-$wave approximation minimally modifies the correlation functions shown in Figs. 6-9. It is rather expected as the correlation functions are significantly different from unity only at small momenta where contributions of higher partial waves are strongly suppressed. The higher partial waves are more important for the functions $S(q_{\rm max})$ but the integrals are still far from being saturated. According to the sum rule (8), the function $S(q_{\rm max})$ should tend to zero for non-identical spheres and to $\pi^3\mathcal{D}_r(0) = (\sqrt{\pi}/2r_0)^3$ for identical ones when $q_{\rm max} \to \infty$. Such a behavior is not observed in our calculations and taking into account the partial waves of higher l does not improve the situation. We see two possible explanations of the problem. 1) The sum rule is in principle fulfilled but one should go to much larger momenta $q_{\rm max}$ to saturate the integral (11). However, higher relative momenta require taking into account more and more partial waves. If this is indeed the case, the sum rule is formally correct but useless because Eq. (1), which is a starting point of the sum rule derivation, assumes that the relative momentum of particles q is much smaller than the typical particle momentum. 2) The integral (11) is divergent as $q_{\rm max}$ goes to infinity. Then, the sum rule is simply meaningless. We discuss the second possibility in more detail in the next section where the Coulomb interaction is studied.

VII. COULOMB INTERACTION

As well known, the Coulomb problem is exactly solvable within the non-relativistic quantum mechanics [14]. The exact wave function of two non-identical particles interacting due to the repulsive Coulomb force is given as

$$\phi_{\mathbf{q}}(\mathbf{r}) = e^{-\frac{\pi\lambda}{2q}}\,\Gamma(1+i\frac{\lambda}{q})\,e^{iqz/2}\,F(-i\frac{\lambda}{q},1,iq\eta)\,, \tag{23}$$

where $\lambda \equiv \mu e^2/8\pi$ with μ being the reduced mass of the two particles and $\pm e$ is the charge of each of them; F denotes the hypergeometric confluent function, and η is the parabolic coordinate (see below). The wave function for the attractive interaction is obtained from (23) by means of the substitution $\lambda \to -\lambda$. When one deals with identical particles, the wave function $\phi_{\mathbf{q}}(\mathbf{r})$ should be replaced by its (anti-)symmetrized form. The modulus of the wave function (23) equals

$$|\phi_{\mathbf{q}}(\mathbf{r})|^2 = G(q)\,|F(-i\frac{\lambda}{q},1,iq\eta)|^2\,,$$

where $G(q)$ is the so-called Gamov factor defined as

$$G(q) = \frac{2\pi\lambda}{q}\,\frac{1}{\exp\left(\frac{2\pi\lambda}{q}\right)-1}\,. \tag{24}$$

As seen, the modulus of the wave function of non-identical particles solely depends on the parabolic coordinate η. Therefore, it is natural to calculate the Coulomb correlation function in the parabolic coordinates: $\eta \equiv r-z$, $\xi \equiv r+z$ and ϕ which is the azimuthal angle. Then, the correlation function computed for the Gaussian source function (4) equals

$$R(q) = \frac{G(q)}{2\sqrt{\pi}\,r_0}\int_0^\infty d\eta\,\exp\Big(-\frac{\eta^2}{16r_0^2}\Big)\,|F(-i\frac{\lambda}{q},1,iq\eta)|^2\,, \tag{25}$$

where the integration over ξ has been performed. It should be noted here that, in contrast to the neutron-proton and hard sphere cases, the Coulomb correlation function is calculated with the *exact* wave function not with the asymptotic form of it.

The modulus of the (anti-)symmetrized Coulomb wave function equals

$$|\phi_{\mathbf{q}}(\mathbf{r})|^2 = \frac{1}{2}\,G(q)\,\Big[|F(-i\frac{\lambda}{q},1,iq\eta)|^2 + |F(-i\frac{\lambda}{q},1,iq\xi)|^2 \pm 2{\rm Re}\Big(e^{iqz}\,F(-i\frac{\lambda}{q},1,iq\eta)\,F^*(-i\frac{\lambda}{q},1,iq\xi)\Big)\Big]\,,$$

and the correlation function analogous to (25) is

$$R(q) = \frac{G(q)}{2\sqrt{\pi}\, r_0}\Big[\int_0^\infty d\eta\, \exp\Big(-\frac{\eta^2}{16 r_0^2}\Big)\, |F(-i\frac{\lambda}{q}, 1, iq\eta)|^2 \tag{26}$$

$$\pm \frac{1}{4r_0^2}\int_0^\infty d\eta \int_0^\infty d\xi\, (\eta+\xi)\, \exp\Big(-\frac{(\eta+\xi)^2}{16 r_0^2}\Big)\, \mathrm{Re}\Big(e^{iq(\xi-\eta)/2}\, F(-i\frac{\lambda}{q}, 1, iq\eta)\, F^*(-i\frac{\lambda}{q}, 1, iq\xi)\Big)\Big]\,.$$

In Figs. 14 and 15 we demonstrate the correlation function of non-identical particles and of identical pions, respectively, which are given by Eqs. (25,26). In the case of non-identical particles, we have taken the reduced mass, which enters the parameter λ, to be equal to that of identical pions. Therefore, the two systems differ only due to the effects of quantum statistics. In both cases the Coulomb interaction is repulsive. While the correlation of non-identical particles is everywhere negative, the correlation of identical pions is negative at small q due to the Coulomb repulsion but it is positive for larger q due to the effect of quantum statistics.

Figs. 16, 17, and 18 show the functions $S(q_{\rm max})$ defined by Eq. (11) which are computed for the correlation functions presented in Figs. 14 and 15, respectively. According to the sum rule (8), $S(q_{\rm max})$ should vanish for sufficiently large $q_{\rm max}$ in the case of non-identical particles which repel each other, and $S(q_{\rm max})$ should tend to $\pi^3\, \mathcal{D}_r(0)$ for identical pions. As seen in Figs. 16, 17, and 18, the functions $S(q_{\rm max})$ do not seem to saturate at large $q_{\rm max}$ and the sum rule is badly violated.

What is wrong here? The derivation of the sum rule (8) implicitly assumes that the integral in the l.h.s. exists *i.e.* it is convergent. Otherwise interchanging of the integrations over $\mathbf{q}$ and over $\mathbf{r}$, which leads to Eq. (5), is mathematically illegal. Unfortunately, the Coulomb correlation functions appear to decay too slowly with q, and consequently the integral in the left hand side of Eq. (8) diverges.

To clarify how the integral diverges, one has to find the asymptotics of the correlation function at large q. This is a difficult problem as one has to determine a behavior of the wave function (23) at large q for any $\mathbf{r}$, and then to perform the integration over $\mathbf{r}$. Up to our knowledge, the problem of large q asymptotics of the Coulomb correlation function has not been satisfactory solved although it has been discussed in several papers [15–19]. We have not found a complete solution of the problem but our rather tedious analysis, which uses the analytic approximate expressions of the hypergeometric confluent function at small and large distances, suggests, in agreement with [15–19], that $(R(q)-1) \sim 1/q^2$ when $q \to \infty$. Then, the integral in the left hand side of Eq. (8) linearly diverges, and the sum rule does not hold. We note that the Gamov factor (24), which represents a zero size source and decays as $1/q$ at large q, leads to the quadratic divergence of the integral (8). We also note here that the asymptotics $1/q^2$ of the correlation function does not have much to do with the well known classical limit of the correlation function [15–19]. Since the large q limit of the correlation function corresponds to the small separation of the charged particles, which at sufficiently large q is smaller than the de Broglie wave length, the classical approximation breaks down.

VIII. FINAL REMARKS

The sum rule (8) provides a rigorous constraint on the correlation function if the momentum integral in Eq. (8) exists. The rule is trivially satisfied by the correlation function of non-interaction particles. The model calculations of the n-p correlation function in the $s-$wave approximation fail to fulfill the sum rule. One suspects that the approximation, which is sufficient to properly describe a general shape of the correlation function, distorts its tail. Since even small deviations of the correlation function from unity at large q generate a sizable contribution to the sum-rule integral, it seems reasonable to expect that the higher partial waves have to be included to comply with the sum rule. Paradoxically, when the higher partial waves are taken into account for the interacting hard spheres the situation is not improved but it gets worse. This suggests that either one should go to really large momenta to saturate the sum-rule integral or the integral in Eq. (8) is divergent. In the case of Coulomb repelling interaction, we certainly deal with the second option. Due to the strong electrostatic repulsion at small distances, the correlation function decays too slowly at large momenta, and consequently the sum-rule integral does not exist.

Being rather useless, the sum rule explains some qualitative features of the correlation function. In particular, it shows that, in spite of the attractive interaction, the correlation can be negative as it happens in the triplet state of neutron-proton pair and in the $\bar{p}\Lambda$ system.

ACKNOWLEDGMENTS

We are very grateful to Wojciech Florkowski for a generous help in performing numerical calculations, and to David Brown, Cristina Manuel, Richard Lednický, and Scott Pratt for fruitful discussions and correspondence.

[1] D. H. Boal, C. K. Gelbke, and B. K. Jennings, Rev. Mod. Phys. **62**, 553 (1990).
[2] U. W. Heinz and B. V. Jacak, Ann. Rev. Nucl. Part. Sci. **49**, 529 (1999).
[3] St. Mrówczyński, Phys. Lett. B **345**, 393 (1995).
[4] G. F. Bertsch, Phys. Rev. Lett. **72**, 2349 (1994) [Erratum-ibid. **77**, 789 (1996)].
[5] D. A. Brown, S. Y. Panitkin, and G. Bertsch, Phys. Rev. C **62**, 014904 (2000) [arXiv:nucl-th/0002039].
[6] D. A. Brown and P. Danielewicz, Phys. Lett. B **398**, 252 (1997) [arXiv:nucl-th/9701010].
[7] D. A. Brown and P. Danielewicz, Phys. Rev. C **64**, 014902 (2001) [arXiv:nucl-th/0010108].
[8] A. Kisiel [STAR Collaboration], J. Phys. G **30**, S1059 (2004) [arXiv:nucl-ex/0403042].
[9] M. I. Podgoretsky, Sov. J. Nucl. Phys. **54**, 891 (1991) [Yad. Fiz. **54**, 1461 (1991)].
[10] E. Janke, F. Emde, and F. Lösch, *Tafeln Höherer Funktionen* (Teubner Verlagsgesellschaft, Stuttgart, 1960).
[11] R. Lednicky and V.L. Lyuboshitz, Sov. J. Nucl. Phys. **35**, 770 (1982) [Yad. Fiz. **35**, 1316 (1982)].
[12] I.E. McCarthy, *Introduction to Nuclear Theory* (John Wiley & Sons, New York, 1968).
[13] P.E. Hodgson, *Nuclear Reactions and Nuclear Structure* (Clarendon Press, Oxford, 1971).
[14] L.I. Schiff, *Quantum Mechanics* (McGraw-Hill, New York, 1968).
[15] M. Gyulassy and S. K. Kauffmann, Nucl. Phys. A **362**, 503 (1981).
[16] Y.D. Kim *et al.*, Phys. Rev. C **45**, 387 (1992).
[17] G. Baym and P. Braun-Munzinger, Nucl. Phys. A **610**, 286C (1996).
[18] Y. Sinyukov, R. Lednicky, S. V. Akkelin, J. Pluta, and B. Erazmus, Phys. Lett. B **432**, 248 (1998).
[19] S. Pratt and S. Petriconi, Phys. Rev. C **68**, 054901 (2003).



FIG. 1. The singlet correlation function of neutron and proton computed for three values of the source size parameter r_0.



FIG. 2. The triplet correlation function of neutron and proton computed for three values of the source size parameter r_0.



FIG. 3. The deuteron formation rate as a function of the source size parameter r_0.



FIG. 4. The function $S(q_{\max})$ for to the singlet correlation function of neutron and proton for three values of the source size parameter r_0. To set the scale the corresponding values of the deuteron formation rate are given.



FIG. 5. The function $S(q_{\max})$ for to the triplet correlation function of neutron and proton for three values of the source size parameter r_0.



FIG. 6. The correlation function of non-identical hard spheres for the source size parameter $r_0 = 4$ fm.



FIG. 7. The correlation function of non-identical hard spheres for the source size parameter $r_0 = 6$ fm.



FIG. 8. The correlation function of identical hard spheres for the source size parameter $r_0 = 4$ fm.



FIG. 9. The correlation function of identical hard spheres for the source size parameter $r_0 = 6$ fm.



FIG. 10. The function $S(q_{\max})$ corresponding to the correlation function of non-identical hard spheres for the source size parameter $r_0 = 4$ fm.



FIG. 11. The function $S(q_{\max})$ corresponding to the correlation function of non-identical hard spheres for the source size parameter $r_0 = 6$ fm.



FIG. 12. The function $S(q_{\max})$ corresponding to the correlation function of identical hard spheres for the source size parameter $r_0 = 4$ fm.



FIG. 13. The function $S(q_{\max})$ corresponding to the correlation function of identical hard spheres for the source size parameter $r_0 = 6$ fm.



FIG. 14. The Coulomb correlation function of non-identical repelling particles for three values of the source size parameter r_0.



FIG. 15. The Coulomb correlation function of identical pions for three values of the source size parameter r_0.



FIG. 16. The function $S(q_{\rm max})$ corresponding to the Coulomb correlation function of non-identical repelling particles for three values of the source size parameter r_0.



FIG. 17. The function $S(q_{\rm max})$ corresponding to the Coulomb correlation function of identical pions for the source size parameter $r_0 = 1$ fm. The dotted line represents the value of $\pi^3\,\mathcal{D}_r(0)$.



FIG. 18. The function $S(q_{\max})$ corresponding to the Coulomb correlation function of identical pions for the source size parameter r_0 equal 3 fm and 5 fm. The dotted lines represents the respective values of $\pi^3\,\mathcal{D}_r(0)$.